

When to Get Baby Shots

Age Group	Hep B hepatitis B	DTP diphtheria, tetanus, pertussis (whooping cough)	Polio	MMR measles, mumps, rubella	Hib <i>Haemophilus influenzae</i> , type b	Varicella (chickenpox)	PCV pneumococcal vaccine	Hep A hepatitis A	Rotavirus	Influenza (flu)
Birth	✓									
2 months	✓	✓	✓		✓		✓		✓	
4 months	1-2 months after the first hep B shot	✓	✓		✓		✓		✓	
6 months		✓			✓		✓		✓	
12-18 months	✓ 6-18 months	✓ 15-18 months	✓ 6-18 months	✓ 12-15 months	✓ 12-15 months	✓ 12-18 months	✓ 12-15 months	✓✓ 2 shots at least 6 months apart		✓ Each fall*
State law requires these shots for enrollment in:	Grades K&7	Child care Grades K-12	Child care Grades K-12	Child care Grades K-12 (Must be given on or after child's first birthday)	Child care	Child care Grades K&7 (Must be given on or after child's first birthday)	Child care, ages 2-24 months			

This schedule may vary a bit, depending on your specific doctor and your child.

***Influenza:** All children age 6 months through 18 years should get a yearly flu vaccination – especially those with risk factors like asthma and diabetes. If a child is younger than 9 years old and is receiving flu vaccine for the first time, they need 2 shots at least 4 weeks apart.

Minnesota law requires written proof of certain shots for children in licensed child care. However, if a child has a medical reason or if his/her parents are conscientiously opposed to any or all of the shots, a legal exemption is available. To file an exemption you must fill out a form, which you can get at child care centers and schools.



Immunization Program
P.O. Box 64975
St. Paul, MN 55164-0975
651-201-5503, 800-657-3970
www.health.state.mn.us/immunize

IC#141-0398 (3/2008)